

PERIODIC APPROXIMATION OF LYAPUNOV EXPONENTS FOR BANACH COCYCLES

BORIS KALININ AND VICTORIA SADOVSKAYA*

ABSTRACT. We consider group-valued cocycles over dynamical systems. The base system is a homeomorphism f of a metric space satisfying a closing property, for example a hyperbolic dynamical system or a subshift of finite type. The cocycle \mathcal{A} takes values in the group of invertible bounded linear operators on a Banach space and is Hölder continuous. We prove that upper and lower Lyapunov exponents of \mathcal{A} with respect to an ergodic invariant measure μ can be approximated in terms of the norms of the values of \mathcal{A} on periodic orbits of f . We also show that these exponents cannot always be approximated by the exponents of \mathcal{A} with respect to measures on periodic orbits. Our arguments include a result of independent interest on construction and properties of a Lyapunov norm for infinite dimensional setting. As a corollary, we obtain estimates of the growth of the norm and of the quasiconformal distortion of the cocycle in terms of the growth at the periodic points of f .

1. INTRODUCTION AND STATEMENTS OF THE RESULTS

Cocycles with values in a group of linear operators on a vector space V are the prime examples of non-commutative cocycles over dynamical systems. For finite dimensional V they have been extensively studied, with examples including random matrices and derivative cocycles of smooth dynamical systems.

The infinite dimensional case is more difficult and so far is much less developed. The simplest examples are given by random and Markov sequences of operators. They correspond to locally constant cocycles over subshifts of finite type, which can be viewed as symbolic systems with hyperbolic behavior. Similarly to finite dimensional case, the derivative of a smooth infinite dimensional system gives a natural example of an operator valued cocycle. We refer to monograph [LL] for an overview of results in this area and to [GoKa, LY12, M12] for some of the recent developments.

In this paper we consider a general setting of cocycles of bounded operators on a Banach space and focus on Hölder continuous cocycles over dynamical systems with hyperbolic behavior. Some of our results hold for measurable cocycles over measure preserving systems, which are defined similarly.

Date: July 23, 2016.

Mathematical subject classification: 37H15, 37D20, 37D25.

Keywords: Cocycles, Lyapunov exponents, periodic orbits, hyperbolic systems.

* Supported in part by NSF grant DMS-1301693.

Definition 1.1. Let f be a homeomorphism of a metric space X , let G be a topological group equipped with a complete metric d , and let A be a function from X to G . The G -valued cocycle over f generated by A is the map $\mathcal{A} : X \times \mathbb{Z} \rightarrow G$ defined by

$$\begin{aligned} \mathcal{A}(x, 0) &= e_G, & \mathcal{A}(x, n) &= \mathcal{A}_x^n = A(f^{n-1}x) \circ \cdots \circ A(x), \\ \mathcal{A}(x, -n) &= \mathcal{A}_x^{-n} = (\mathcal{A}_{f^{-n}x}^n)^{-1}, & n &\in \mathbb{N}. \end{aligned}$$

Clearly, \mathcal{A} satisfies the cocycle equation $\mathcal{A}_x^{n+k} = \mathcal{A}_{f^kx}^n \circ \mathcal{A}_x^k$.

We say that the cocycle \mathcal{A} is bounded if its generator A is a bounded function and that \mathcal{A} is α -Hölder if A is Hölder continuous with exponent $0 < \alpha \leq 1$ with respect to the metric d , i.e. there exists $M > 0$ such that

$$(1.1) \quad d(A(x), A(y)) \leq M \operatorname{dist}(x, y)^\alpha \quad \text{for all sufficiently close } x, y \in X.$$

In this paper we consider cocycles with values in the group of invertible operators on a Banach space V . The space $L(V)$ of bounded linear operators on V is a Banach space equipped with the operator norm $\|A\| = \sup\{\|Av\| : v \in V, \|v\| \leq 1\}$. The open set $GL(V)$ of invertible elements in $L(V)$ is a topological group and a complete metric space with respect to the metric

$$d(A, B) = \|A - B\| + \|A^{-1} - B^{-1}\|.$$

We call a $GL(V)$ -valued cocycle \mathcal{A} a *Banach cocycle*.

Definition 1.2. Let μ be an ergodic f -invariant Borel probability measure on X . The upper and lower Lyapunov exponents of \mathcal{A} with respect to μ are

$$(1.2) \quad \begin{aligned} \lambda(\mathcal{A}, \mu) &= \lim_{n \rightarrow \infty} \frac{1}{n} \log \|\mathcal{A}_x^n\| \quad \text{for } \mu \text{ almost every } x \in X, \\ \chi(\mathcal{A}, \mu) &= \lim_{n \rightarrow \infty} \frac{1}{n} \log \|(\mathcal{A}_x^n)^{-1}\|^{-1} \quad \text{for } \mu \text{ almost every } x \in X. \end{aligned}$$

Each of these limits exists and is the same μ almost everywhere by the Subadditive Ergodic Theorem, see Section 2. If V is finite dimensional, these are precisely the largest and smallest of the Lyapunov exponents given by the Multiplicative Ergodic Theorem of Oseledets.

Our goal is to obtain a periodic approximation for the exponents of a cocycle over a system with hyperbolic behavior. To streamline the arguments we formulate explicitly the property that we will use.

Definition 1.3. We say that a homeomorphism f of a metric space X satisfies the closing property if there exist constants $D, \gamma, \delta_0 > 0$ such that for any $x \in X$ and $k \in \mathbb{N}$ with $\operatorname{dist}(x, f^kx) < \delta_0$ there exists a point $p \in X$ with $f^kp = p$ such that the orbit segments x, fx, \dots, f^kx and p, fp, \dots, f^kp are exponentially close, more precisely,

$$(1.3) \quad \operatorname{dist}(f^i x, f^i p) \leq D \operatorname{dist}(x, f^k x) e^{-\gamma \min\{i, k-i\}} \quad \text{for every } i = 0, \dots, k.$$

Systems satisfying this closing property include symbolic dynamical systems such as subshifts of finite type as well as smooth hyperbolic systems such as hyperbolic automorphisms of tori and nilmanifolds, Anosov diffeomorphisms, and hyperbolic maps of locally maximal sets. For the smooth systems the closing property follows from Anosov Closing Lemma, see e.g. [KtH, 6.4.15-17].

Now we state our main result. Note that we do not assume compactness of X .

Theorem 1.4. *Let X be separable metric space, let f be a homeomorphism of X satisfying the closing property, let μ be an ergodic f -invariant Borel probability measure on X , and let \mathcal{A} be a bounded Hölder continuous Banach cocycle over f . Then for each $\epsilon > 0$ there exists a periodic point $p = f^k p$ in X such that*

$$(1.4) \quad \left| \lambda(\mathcal{A}, \mu) - \frac{1}{k} \log \|\mathcal{A}_p^k\| \right| < \epsilon \quad \text{and} \quad \left| \chi(\mathcal{A}, \mu) - \frac{1}{k} \log \|(\mathcal{A}_p^k)^{-1}\|^{-1} \right| < \epsilon.$$

Moreover, for any $N \in \mathbb{N}$ there exists such $p = f^k p$ with $k > N$.

A periodic approximation of Lyapunov exponents for finite-dimensional V was established in [K11, Theorem 1.4] under slightly stronger closing assumption. It was shown, in particular, that for each $\epsilon > 0$ there exists a point $p = f^k p$ such that for the uniform measure μ_p on its orbit

$$(1.5) \quad |\lambda(\mathcal{A}, \mu) - \lambda(\mathcal{A}, \mu_p)| < \epsilon \quad \text{and} \quad |\chi(\mathcal{A}, \mu) - \chi(\mathcal{A}, \mu_p)| < \epsilon,$$

and it is clear from the argument that (1.4) also holds. We note that

$$\lambda(\mathcal{A}, \mu_p) = (1/k) \log (\text{spectral radius of } \mathcal{A}_p^k)$$

and so in our setting for $p = f^k p$ satisfying (1.4) we have

$$(1.6) \quad \lambda(\mathcal{A}, \mu_p) < \lambda(\mathcal{A}, \mu) + \epsilon \quad \text{and} \quad \chi(\mathcal{A}, \mu_p) > \chi(\mathcal{A}, \mu) - \epsilon.$$

However, approximation (1.5) is not always possible in infinite-dimensional setting. The following proposition is based on an example by L. Gurvits of a pair of operators whose joint spectral radius is greater than the generalized spectral radius [Gu].

Proposition 1.5. *There exists a locally constant cocycle \mathcal{A} over a full shift on two symbols and an ergodic invariant measure μ such that $\lambda(\mathcal{A}, \mu) > \sup_{\mu_p} \lambda(\mathcal{A}, \mu_p)$, where the supremum is taken over all uniform measures μ_p on periodic orbits.*

The proof of the finite dimensional periodic approximation result in [K11] relies on Multiplicative Ergodic Theorem, which yields that the cocycle has finitely many Lyapunov exponents and, in particular, the largest one is isolated. As this may not be the case in infinite dimensional setting even for a single operator, we have to use a different approach. First we construct a measurable Lyapunov norm for infinite dimensional setting and establish its temperedness. This is a result of independent interest. We use this norm to obtain estimates of the growth of the cocycle for trajectories close to ones with regular behavior. We also use further developments of the subadditive ergodic theorem obtained in [KaM99, GoKa] and extend various

techniques from [K11] and [G]. Our methods may also be useful in the study of cocycles with values in diffeomorphism groups as well as of infinite-dimensional non-uniformly hyperbolic dynamical systems on Hilbert or Banach manifolds.

As a corollary of our main result, we obtain estimates of the growth of the norm and of the *quasiconformal distortion* $Q(x, n) := \|\mathcal{A}_x^n\| \cdot \|(\mathcal{A}_x^n)^{-1}\|$ of a cocycle \mathcal{A} in terms of the growth at periodic points.

Corollary 1.6. *Let f be a homeomorphism of a compact metric space X satisfying the closing property and let \mathcal{A} be a Hölder continuous Banach cocycle over f . Then*

$$(i) \quad \limsup_{n \rightarrow \infty} \{\|\mathcal{A}_x^n\|^{1/n} : x \in X\} = \limsup_{k \rightarrow \infty} \sup \{\|\mathcal{A}_p^k\|^{1/k} : p = f^k p \in X\}.$$

In particular, if for some numbers C and s we have $\|\mathcal{A}_p^k\| \leq Ce^{sk}$ whenever $p = f^k p$, then for each $\epsilon > 0$ there exists a number C_ϵ such that

$$\|\mathcal{A}_x^n\| \leq C_\epsilon e^{(s+\epsilon)n} \quad \text{for all } x \in X \text{ and } n \in \mathbb{N}.$$

$$(ii) \quad \limsup_{n \rightarrow \pm\infty} \{Q(x, n)^{1/|n|} : x \in X\} = \limsup_{k \rightarrow \infty} \sup \{Q(p, k)^{1/k} : p = f^k p \in X\}.$$

In particular, if for some numbers C and s we have $Q(p, k) \leq Ce^{sk}$ whenever $p = f^k p$, then for each $\epsilon > 0$ there exists a number C'_ϵ such that

$$Q(x, n) \leq C'_\epsilon e^{(s+\epsilon)|n|} \quad \text{for all } x \in X \text{ and } n \in \mathbb{Z}.$$

The number $\hat{\lambda}(\mathcal{A}) = \lim_{n \rightarrow \infty} \sup \{\|\mathcal{A}_x^n\|^{1/n} : x \in X\}$ gives the maximal growth rate of the cocycle and it is known that $\hat{\lambda}(\mathcal{A}) = \sup_\mu \lambda(\mathcal{A}, \mu)$, where the supremum is taken over all ergodic f -invariant measures. Part (i) of the corollary can also be deduced from a recent work of M. Guysinsky [G], where he showed that for each $\epsilon > 0$ there exists a periodic point $p = f^k p$ such that $\frac{1}{k} \log \|\mathcal{A}_p^k\| > \hat{\lambda}(\mathcal{A}) - \epsilon$. He used this result to obtain a generalization of Livsic periodic point theorem for Banach cocycles.

Part (ii) of the corollary has a broad scope of applications as the quasiconformal distortion $Q(x, n)$ is used in so called *fiber bunching* condition for cocycles over hyperbolic systems. This condition means that $Q(x, n)$ is dominated by expansion and contraction in the base system and it plays a crucial role in most of the results on non-commutative cocycles over hyperbolic systems. The corollary allows to obtain fiber bunching from the periodic data.

Remark 1.7. *More generally, Theorem 1.4 and Corollary 1.6 hold if we replace $X \times V$ by a Hölder continuous vector bundle \mathcal{V} over X with fiber V and the cocycle \mathcal{A} by an automorphism $\mathcal{F} : \mathcal{V} \rightarrow \mathcal{V}$ covering f . This setting is described in detail in Section 2.2 of [KS13] and the proofs work without any significant modifications.*

We discuss preliminaries on subadditive cocycles in Section 2, give construction and properties of the Lyapunov norm in Section 3, prove Theorem 1.4 in Section 4, and prove Proposition 1.5 and Corollary 1.6 in Section 5.

2. SUBADDITIVE COCYCLES AND THEIR EXPONENTS

A *subadditive cocycle* over a dynamical system (X, f) is a sequence of functions $a_n : X \rightarrow \mathbb{R}$ such that

$$a_{n+k}(x) \leq a_k(x) + a_n(f^k x) \quad \text{for all } x \in X \text{ and } k, n \in \mathbb{N}.$$

We define $\mathcal{V} = X \times V$ and view \mathcal{A}_x^n as a map from \mathcal{V}_x to $\mathcal{V}_{f^n x}$. While this is not necessary for the trivial bundle, it makes notations and arguments more intuitive and facilitates extension to the bundle setting. We denote

$$(2.1) \quad a_n(x) = \log \|\mathcal{A}_x^n\|, \quad b_n(x) = a_n(f^{-n}x) = \log \|\mathcal{A}_{f^{-n}x}^n\|,$$

$$(2.2) \quad \tilde{a}_n(x) = \log \|(\mathcal{A}_x^n)^{-1}\|, \quad \tilde{b}_n(x) = \tilde{a}_n(f^{-n}x) = \log \|(\mathcal{A}_{f^{-n}x}^n)^{-1}\| = \log \|\mathcal{A}_x^{-n}\|.$$

It is easy to see that $a_n(x)$ and $\tilde{a}_n(x)$ are subadditive cocycles over f and that $b_n(x)$ and $\tilde{b}_n(x)$ are subadditive cocycles over f^{-1} .

For any ergodic measure-preserving transformation f of a probability space (X, μ) and any subadditive cocycle over f with integrable a_n , the Subadditive Ergodic Theorem yields that for μ almost all x

$$\lim_{n \rightarrow \infty} \frac{1}{n} a_n(x) = \lim_{n \rightarrow \infty} \frac{1}{n} a_n(\mu) = \inf_{n \in \mathbb{N}} \frac{1}{n} a_n(\mu) =: \nu(a, \mu), \quad \text{where } a_n(\mu) = \int_X a_n(x) d\mu.$$

The limit $\nu(a, \mu) \geq -\infty$ is called the *exponent* of the cocycle a_n with respect to μ .

With our choice of $a_n(x)$, this theorem gives the existence of the limit in the first equation of (1.2) and that $\nu(a, \mu)$ is the upper Lyapunov exponent $\lambda = \lambda(\mathcal{A}, \mu)$ of the cocycle \mathcal{A} , so for μ almost all x

$$(2.3) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \log \|\mathcal{A}_x^n\| = \lim_{n \rightarrow \infty} \frac{1}{n} a_n(x) = \lambda := \lambda(\mathcal{A}, \mu).$$

Since $b_n(\mu) = \int_X b_n(x) d\mu = \int_X a_n(x) d\mu = a_n(\mu)$, it follows that for μ almost all x

$$(2.4) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \log \|\mathcal{A}_{f^{-n}x}^n\| = \lim_{n \rightarrow \infty} \frac{1}{n} b_n(x) = \lambda.$$

Similarly, for μ almost all x

$$(2.5) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \log \|\mathcal{A}_x^{-n}\| = \lim_{n \rightarrow \infty} \frac{1}{n} \tilde{b}_n(x) = \lim_{n \rightarrow \infty} \frac{1}{n} \tilde{a}_n(x) = -\chi,$$

where $\chi := \chi(\mathcal{A}, \mu)$ is the lower Lyapunov exponent of the cocycle \mathcal{A} defined in (1.2), and it is easy to see that $\chi \leq \lambda$ and both are finite. We denote

$$(2.6) \quad \Lambda = \Lambda_\mu = \{x \in X : \text{equations (2.3), (2.4), and (2.5) hold}\}$$

and in particular both equalities in (1.2) hold. Clearly, $\mu(\Lambda) = 1$.

3. LYAPUNOV NORM

In this section we construct a certain version of Lyapunov or adapted norm for our setting, which allows us to control the norms of \mathcal{A}_x and $(\mathcal{A}_x)^{-1}$. It is cruder than the usual Lyapunov norm for matrix cocycles constructed using Oseledets splitting, which is unavailable our setting. Our construction is closer to that of an adapted metric for an Anosov system. Since the Lyapunov norm in general depends only measurably on x , it is important to provide a comparison with the standard norm. We do this in the second part of Proposition 3.1 below.

For a fixed $\epsilon > 0$ and a point $x \in \Lambda$, the ϵ -Lyapunov norm $\|\cdot\|_x = \|\cdot\|_{x,\epsilon}$ on \mathcal{V}_x is defined as follows. For $u \in \mathcal{V}_x$,

$$(3.1) \quad \|u\|_x = \|u\|_{x,\epsilon} = \sum_{n=0}^{\infty} \|\mathcal{A}_x^n(u)\| e^{-(\lambda+\epsilon)n} + \sum_{n=1}^{\infty} \|\mathcal{A}_x^{-n}(u)\| e^{(\chi-\epsilon)n}.$$

By the definition (2.6) of Λ , both series converge exponentially.

We denote the operator norm with respect to the Lyapunov norms by $\|\cdot\|_{y \leftarrow x}$. For any points $x, y \in \Lambda$ and any linear map $A : \mathcal{V}_x \rightarrow \mathcal{V}_y$ it is defined by

$$\|A\|_{y \leftarrow x} = \sup \{ \|A(u)\|_{y,\epsilon} : u \in \mathcal{V}_x, \|u\|_{x,\epsilon} = 1 \}.$$

Proposition 3.1. *Let f be an ergodic invertible measure-preserving transformation of a probability space (X, μ) and let \mathcal{A} be a bounded measurable Banach cocycle over f with the upper Lyapunov exponent λ and the lower Lyapunov exponent χ . Then for each $\epsilon > 0$ the Lyapunov norm $\|\cdot\|_{x,\epsilon}$ given by (3.1) satisfies the following.*

(i) For each point x in Λ ,

$$(3.2) \quad \|\mathcal{A}_x\|_{f_x \leftarrow x} \leq e^{\lambda+\epsilon} \quad \text{and} \quad \|\mathcal{A}_x^{-1}\|_{f^{-1}x \leftarrow x} \leq e^{-\chi+\epsilon}.$$

(ii) There exists an f -invariant set $\mathcal{R} \subset \Lambda$ with $\mu(\mathcal{R}) = 1$ so that for each $\rho > 0$ there exists a measurable function $K_\rho(x)$ such that for all $x \in \mathcal{R}$

$$(3.3) \quad \|u\| \leq \|u\|_{x,\epsilon} \leq K_\rho(x) \|u\| \quad \text{for all } u \in \mathcal{V}_x, \quad \text{and}$$

$$(3.4) \quad K_\rho(x) e^{-\rho|n|} \leq K_\rho(f^n x) \leq K_\rho(x) e^{\rho|n|} \quad \text{for all } n \in \mathbb{Z}.$$

Proof. (i) Let $u \in \mathcal{V}_x$. Using definition (3.1) and the fact that $\chi \leq \lambda$ we obtain

$$\begin{aligned} \|\mathcal{A}_x(u)\|_{f_x} &= \sum_{n=0}^{\infty} \|\mathcal{A}_{f_x}^n(\mathcal{A}_x(u))\| e^{-(\lambda+\epsilon)n} + \sum_{n=1}^{\infty} \|\mathcal{A}_{f_x}^{-n}(\mathcal{A}_x(u))\| e^{(\chi-\epsilon)n} \\ &= \sum_{n=0}^{\infty} \|\mathcal{A}_x^{n+1}(u)\| e^{-(\lambda+\epsilon)n} + \sum_{n=1}^{\infty} \|\mathcal{A}_x^{-n+1}(u)\| e^{(\chi-\epsilon)n} \end{aligned}$$

$$\begin{aligned}
&= e^{(\lambda+\epsilon)} \left(\sum_{k=1}^{\infty} \|\mathcal{A}_x^k(u)\| e^{-(\lambda+\epsilon)k} + \|u\| e^{(\chi-\epsilon)-(\lambda+\epsilon)} + \sum_{n=2}^{\infty} \|\mathcal{A}_x^{-n+1}(u)\| e^{(\chi-\epsilon)n-(\lambda+\epsilon)} \right) \\
&\leq e^{(\lambda+\epsilon)} \left(\sum_{k=1}^{\infty} \|\mathcal{A}_x^k(u)\| e^{-(\lambda+\epsilon)k} + \|u\| + \sum_{n=2}^{\infty} \|\mathcal{A}_x^{-n+1}(u)\| e^{(\chi-\epsilon)(n-1)} \right) \\
&= e^{(\lambda+\epsilon)} \left(\sum_{k=0}^{\infty} \|\mathcal{A}_x^k(u)\| e^{-(\lambda+\epsilon)k} + \sum_{k=1}^{\infty} \|\mathcal{A}_x^{-k}(u)\| e^{(\chi-\epsilon)k} \right) = e^{(\lambda+\epsilon)} \|u\|_x,
\end{aligned}$$

and the first inequality follows. The second inequality is obtained similarly.

(ii) The uniform lower bound $\|u\|_{x,\epsilon} \geq \|u\|$ follows immediately from the definition of $\|u\|_{x,\epsilon}$, so it remains to establish the upper bound.

By (2.6) for each $\epsilon > 0$ and $x \in \Lambda$ there exists $N_\epsilon(x) \in \mathbb{N}$ such that

$$(3.5) \quad \|\mathcal{A}_x^n\| \leq e^{(\lambda+\epsilon)n} \quad \text{and} \quad \|\mathcal{A}_x^{-n}\| \leq e^{(-\chi+\epsilon)n} \quad \text{for all } n > N_\epsilon(x).$$

Then for all $n \in \mathbb{N}$

$$(3.6) \quad \|\mathcal{A}_x^n\| \leq M_\epsilon(x) e^{(\lambda+\epsilon)n} \quad \text{and} \quad \|\mathcal{A}_x^{-n}\| \leq M'_\epsilon(x) e^{(-\chi+\epsilon)n}, \quad \text{where}$$

$$(3.7) \quad \begin{aligned} M_\epsilon(x) &= \max \{ \|\mathcal{A}_x^n\| e^{-(\lambda+\epsilon)n} : 0 \leq n \leq N_\epsilon(x) \} \quad \text{and} \\ M'_\epsilon(x) &= \max \{ \|\mathcal{A}_x^{-n}\| e^{(\chi-\epsilon)n} : 0 \leq n \leq N_\epsilon(x) \}. \end{aligned}$$

We note that $M_\epsilon(x) \geq \|\mathcal{A}_x^0\| = 1$ and, since $\|\mathcal{A}_x^n\| e^{-(\lambda+\epsilon)n} \leq 1$ for all $n > N_\epsilon$,

$$M_\epsilon(x) = \sup \{ \|\mathcal{A}_x^n\| e^{-(\lambda+\epsilon)n} : n \geq 0 \}.$$

It follows, in particular, that the function M_ϵ is measurable. Similarly, M'_ϵ is also measurable. Using (3.6) with $\epsilon/2$, we estimate $\|u\|_{x,\epsilon}$ as follows.

$$\begin{aligned}
\|u\|_{x,\epsilon} &= \sum_{n=0}^{\infty} \|\mathcal{A}_x^n(u)\| e^{-(\lambda+\epsilon)n} + \sum_{n=1}^{\infty} \|\mathcal{A}_x^{-n}(u)\| e^{(\chi-\epsilon)n} \leq \\
&\leq \sum_{n=0}^{\infty} \|u\| M_{\epsilon/2}(x) e^{(\lambda+\epsilon/2)n} e^{-(\lambda+\epsilon)n} + \sum_{n=1}^{\infty} \|u\| M'_{\epsilon/2}(x) e^{(-\chi+\epsilon/2)n} e^{(\chi-\epsilon)n} \\
&= \|u\| \cdot (M_{\epsilon/2}(x) + M'_{\epsilon/2}(x)) / (1 - e^{-\epsilon/2}) =: \|u\| \cdot \tilde{M}_\epsilon(x).
\end{aligned}$$

By Lemma 3.2 below, the function $\tilde{M}_\epsilon(x)$ is tempered on a set \mathcal{R} of full measure. Hence by [BP, Lemma 3.5.7] for each $\rho > 0$,

$$K_\rho(x) = \sum_{n \in \mathbb{Z}} \tilde{M}_\epsilon(f^n x) e^{-\rho|n|}$$

is a measurable function defined on \mathcal{R} satisfying (3.4). The inequality (3.3) follows since $\tilde{M}_\epsilon(x) \leq K_\rho(x)$. \square

Lemma 3.2. *For each $\epsilon > 0$ the functions M_ϵ and M'_ϵ defined by (3.7) are forward and backward tempered on a set $\mathcal{R} \subset \Lambda$ with $\mu(\mathcal{R}) = 1$, that is for all $x \in \mathcal{R}$*

$$(i) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \log M_\epsilon(f^n x) = 0 \quad \text{and} \quad (ii) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \log M_\epsilon(f^{-n} x) = 0$$

and similarly for M'_ϵ .

Proof. We fix $\epsilon > 0$ and give the proof for M_ϵ . The result for M'_ϵ follows by reversing the time. We note that $M_\epsilon \geq 1$ so that the sequences in (i) and (ii) are nonnegative.

(i) First we show that M_ϵ is forward tempered. We consider $s > 0$ and suppose that for some $x \in \Lambda$

$$(3.8) \quad \limsup_{n \rightarrow \infty} \frac{1}{n} \log M_\epsilon(f^n x) > s > 0.$$

Then there exist infinitely many $k \in \mathbb{N}$ such that $M_\epsilon(f^k x) > e^{sk}$, and for any such k there exists $n = n(k)$ such that

$$(3.9) \quad \|\mathcal{A}_{f^k x}^n\| = M_\epsilon(f^k x) e^{(\lambda+\epsilon)n} > e^{sk} e^{(\lambda+\epsilon)n}.$$

Since the cocycle \mathcal{A} is bounded, there exist constants $\lambda' > \lambda$ and $\chi' < \chi$ such that

$$(3.10) \quad \|\mathcal{A}_x\| \leq e^{\lambda'} \quad \text{and} \quad \|\mathcal{A}_x^{-1}\| \leq e^{-\chi'} \quad \text{for all } x \in X.$$

First we show that if k and n satisfy inequalities (3.9) and $k > N_{\epsilon/2}(x)$, where $N_{\epsilon/2}(x)$ is defined as in (3.5), then they must be comparable, more precisely,

$$(3.11) \quad ck < n < Ck, \quad \text{where} \quad C = 1 + 2(\lambda - \chi')/\epsilon \quad \text{and} \quad c = s/(\lambda' - \lambda).$$

Indeed, for such k inequality (3.5) implies $\|\mathcal{A}_x^{n+k}\| \leq e^{(\lambda+\epsilon/2)(n+k)}$ so we obtain

$$\|\mathcal{A}_{f^k x}^n\| = \|\mathcal{A}_x^{n+k} \circ (\mathcal{A}_x^k)^{-1}\| \leq \|\mathcal{A}_x^{n+k}\| \cdot \|(\mathcal{A}_x^k)^{-1}\| \leq e^{(\lambda+\epsilon/2)(n+k)} e^{-\chi'k}.$$

Together with (3.9) this yields $e^{sk} e^{(\lambda+\epsilon)n} \leq e^{(\lambda+\epsilon/2)(n+k)} e^{-\chi'k}$ and so

$$n \leq k \cdot 2(\lambda - \chi' - s + \epsilon/2)/\epsilon < Ck.$$

On the other hand, (3.9) and $\|\mathcal{A}_{f^k x}^n\| \leq e^{\lambda'n}$ imply $e^{sk} e^{(\lambda+\epsilon)n} \leq e^{\lambda'n}$ and so

$$n \geq k \cdot s/(\lambda' - \lambda - \epsilon) > ck.$$

Now we show that there exists a set of full measure such that for each x in this set (3.8) leads to a contradiction. We choose $\delta = \delta(s)$ so that

$$(3.12) \quad 0 < \delta < c/(C+1), \quad \text{where } c \text{ and } C \text{ are as in (3.11)}.$$

By Egorov's theorem, there exists a set $Y = Y(\delta)$ with $\mu(Y) > 1 - \delta$ so that the convergence in (2.4) is uniform on Y , and thus there exists an integer $N'_\epsilon = N'_\epsilon(Y)$ such that

$$(3.13) \quad b_n(y) < (\lambda + \epsilon)n \quad \text{for all } y \in Y \text{ and } n > N'_\epsilon.$$

Let Λ_Y be the set of full μ measure on which Birkhoff Ergodic Theorem holds for the indicator function of Y , i.e. for all $x \in \Lambda_Y$

$$\lim_{N \rightarrow \infty} \frac{1}{N} \#\{\ell : 1 \leq \ell \leq N \text{ and } f^\ell x \in Y\} = \mu(Y) > 1 - \delta.$$

Now we consider $x \in \Lambda \cap \Lambda_Y$. Then for all sufficiently large k we have

$$\#\{\ell : 1 \leq \ell \leq (k+n) \text{ and } f^\ell x \notin Y\} < \delta(k+n).$$

Hence there exists $n' \leq n$ with $n - n' < \delta(k+n)$ such that $y = f^{k+n'}x \in Y$. Using (3.11) we get

$$(3.14) \quad n - n' < \delta(k+n) < \delta(k+Ck) = \delta(C+1)k.$$

Using again (3.11) we obtain

$$n' = n - (n - n') > ck - \delta(k+Ck) = (c - \delta(C+1))k,$$

where $c - \delta(C+1) > 0$ by (3.12). Therefore, since k can be chosen arbitrarily large, we can assume that $n' > N'_\epsilon$. Then, as $y = f^{k+n'}x \in Y$, we can estimate

$$\|\mathcal{A}_{f^k x}^n\| \leq \|\mathcal{A}_{f^k x}^{n'}\| \cdot \|\mathcal{A}_{f^{k+n'} x}^{n-n'}\| \leq e^{b_{n'}(f^{k+n'}x)} e^{\lambda'(n-n')} < e^{(\lambda+\epsilon)n'} e^{\lambda'(n-n')}.$$

Combining this with (3.9), we obtain

$$e^{sk} e^{(\lambda+\epsilon)n} < \|\mathcal{A}_{f^k x}^n\| < e^{(\lambda+\epsilon)n'} e^{\lambda'(n-n')}$$

so that $sk < (\lambda' - \lambda - \epsilon)(n - n')$. Then using (3.14), (3.12), and (3.11) we obtain a contradiction

$$sk < (\lambda' - \lambda - \epsilon)\delta k(C+1) < (\lambda' - \lambda)ck = sk.$$

Thus for each $s > 0$ there exists a full measure set $\Lambda_Y = \Lambda_Y(s)$ such that

$$\limsup \frac{1}{n} \log M_\epsilon(f^n x) \leq s \quad \text{for all } x \in \Lambda \cap \Lambda_Y,$$

and the first part of the lemma follows by taking the intersection of Λ and $\Lambda_Y(1/n)$, $n \in \mathbb{N}$.

(ii) Now we show that M_ϵ is backward tempered using a similar argument. Suppose that for some $x \in \Lambda$

$$\limsup \frac{1}{n} \log M_\epsilon(f^{-n}x) > s > 0.$$

Then $M_\epsilon(f^{-k}x) > e^{sk}$ for infinitely many $k \in \mathbb{N}$, and hence for some $n = n(k)$

$$(3.15) \quad \|\mathcal{A}_{f^{-k}x}^n\| > e^{sk} e^{(\lambda+\epsilon)n}.$$

By (2.4) there exists $N'_\epsilon(x)$ such that $b_k(x) < e^{(\lambda+\epsilon)k}$ for $k > N'_\epsilon(x)$. If $n > k > N'_\epsilon(x)$ then we have

$$\begin{aligned} \|\mathcal{A}_{f^{-k}x}^n\| &\leq \|\mathcal{A}_{f^{-k}x}^k\| \cdot \|\mathcal{A}_x^{n-k}\| \leq e^{b_k(x)} M_\epsilon(x) e^{(\lambda+\epsilon)(n-k)} \\ &\leq M_\epsilon(x) e^{(\lambda+\epsilon)k} e^{(\lambda+\epsilon)(n-k)} = M_\epsilon(x) e^{(\lambda+\epsilon)n}, \end{aligned}$$

which is incompatible with (3.15) for large k , and hence $n \leq k$. Also, (3.15) and $\|\mathcal{A}_{f^{-k}x}^n\| \leq e^{\lambda'n}$ imply $e^{sk}e^{(\lambda+\epsilon)n} \leq e^{\lambda'n}$ and so $n > ck$, where $c = s/(\lambda' - \lambda)$ as in (3.11). We conclude that for large k , any $n = n(k)$ in (3.15) satisfies $ck < n \leq k$.

We choose $\delta = \delta(s)$ so that $0 < \delta < c$ and take sets $Y = Y(\delta)$ with $\mu(Y) > 1 - \delta$ and Λ_Y as in the first part. Then if $x \in \Lambda \cap \Lambda_Y$ and k is sufficiently large there exists $n' \leq n$ with $n - n' < \delta k$ such that $y' = f^{-k+n'}x \in Y$. Since $\delta < c$ and $n' = n - (n - n') > ck - \delta k = (c - \delta)k$ we conclude that $n' > N'_\epsilon$ if k is sufficiently large. Thus for such k we can estimate as before

$$\|\mathcal{A}_{f^{-k}x}^n\| \leq \|\mathcal{A}_{f^{-k}x}^{n'}\| \cdot \|\mathcal{A}_{f^{-k+n'}x}^{n-n'}\| \leq e^{b_{n'}(f^{-k+n'}x)} e^{\lambda'(n-n')} < e^{(\lambda+\epsilon)n'} e^{\lambda'(n-n')}.$$

Combining this with (3.15), we obtain

$$e^{sk}e^{(\lambda+\epsilon)n} < \|\mathcal{A}_{f^{-k}x}^n\| < e^{(\lambda+\epsilon)n'} e^{\lambda'(n-n')},$$

which yields a contradiction as $n - n' < \delta k$ and $\delta < c = s/(\lambda' - \lambda)$:

$$sk < (\lambda' - \lambda - \epsilon)(n - n') < (\lambda' - \lambda)\delta k = sk.$$

Thus on the full measure set $\Lambda \cap \Lambda_Y$ we have $\limsup \frac{1}{n} \log M_\epsilon(f^{-n}x) \leq s$, and the second part of the lemma follows. \square

4. PROOF OF THEOREM 1.4

4.1. Preliminary results. We fix $\epsilon > 0$ and consider the corresponding Lyapunov norm $\|\cdot\|_x = \|\cdot\|_{x,\epsilon}$ given by (3.1). We apply Proposition 3.1 (ii) with $\rho = \epsilon$ and obtain a full measure set $\mathcal{R} \subset \Lambda$ where the function $K = K_\epsilon$ satisfies (3.3) and (3.4). For any $\ell > 1$ we define

$$(4.1) \quad \mathcal{R}_\ell = \{x \in \mathcal{R} : K(x) \leq \ell\},$$

and note that $\mu(\mathcal{R}_\ell) \rightarrow 1$ as $\ell \rightarrow \infty$.

We recall that $\|\mathcal{A}_x^n\|_{x_n \leftarrow x_0} \leq e^{n(\lambda+\epsilon)}$ and $\|(\mathcal{A}_x^n)^{-1}\|_{x_0 \leftarrow x_n} \leq e^{n(-\chi+\epsilon)}$ for $x \in \Lambda$. In Lemma (4.1) we obtain similar estimates for *any* point $y \in X$ whose trajectory is close to that of a point $x \in \mathcal{R}_\ell$. Since the Lyapunov norm may not exist at points $f^n y$ we will use the Lyapunov norms at the corresponding points $f^n x$ for the estimates. Since the bundle \mathcal{V} is trivial this creates no problem. For a non-trivial bundle one would need to identify spaces $\mathcal{V}_{f^n x}$ and $\mathcal{V}_{f^n y}$, which can be done for nearby points.

Lemma 4.1. *Let f be an ergodic invertible measure-preserving transformation of a probability space (X, μ) and let \mathcal{A} be a bounded α -Hölder Banach cocycle over f with the upper Lyapunov exponent λ and the lower Lyapunov exponent χ .*

Then for any $\gamma > \epsilon/\alpha$ there exists a constant $c = c(\mathcal{A}, \alpha\gamma - \epsilon)$ such that for any point x in \mathcal{R}_ℓ with $f^m x$ in \mathcal{R}_ℓ and any point $y \in X$ such that the orbit segments $x, fx, \dots, f^m x$ and $y, fy, \dots, f^m y$ satisfy with some $\delta > 0$

$$(4.2) \quad \text{dist}(f^i x, f^i y) \leq \delta e^{-\gamma \min\{i, m-i\}} \quad \text{for every } i = 0, \dots, m$$

we have for all $0 \leq n \leq m$

$$(4.3) \quad \|\mathcal{A}_y^n\| \leq \ell \|\mathcal{A}_y^n\|_{x_n \leftarrow x_0} \leq \ell e^{c\ell\delta^\alpha} e^{n(\lambda+\epsilon)} \quad \text{and}$$

$$(4.4) \quad \|(\mathcal{A}_y^n)^{-1}\| \leq \ell e^{\epsilon \min\{n, m-n\}} \|(\mathcal{A}_y^n)^{-1}\|_{x_0 \leftarrow x_n} \leq \ell e^{\epsilon \min\{n, m-n\}} e^{c\ell\delta^\alpha} e^{n(-\lambda+\epsilon)}.$$

Proof. First we prove (4.3). We denote

$$x_i = f^i x \quad \text{and} \quad y_i = f^i y, \quad i = 0, \dots, m,$$

and use (3.2) to estimate the Lyapunov norm for $0 < n \leq m$

$$(4.5) \quad \begin{aligned} \|\mathcal{A}_y^n\|_{x_n \leftarrow x_0} &\leq \prod_{i=0}^{n-1} \|\mathcal{A}_{y_i}\|_{x_{i+1} \leftarrow x_i} \leq \prod_{i=0}^{n-1} \|\mathcal{A}_{x_i}\|_{x_{i+1} \leftarrow x_i} \cdot \|(\mathcal{A}_{x_i})^{-1} \circ \mathcal{A}_{y_i}\|_{x_i \leftarrow x_i} \\ &\leq e^{n(\lambda+\epsilon)} \prod_{i=0}^{n-1} \|(\mathcal{A}_{x_i})^{-1} \circ \mathcal{A}_{y_i}\|_{x_i \leftarrow x_i}. \end{aligned}$$

We consider $\Delta_i = (\mathcal{A}_{x_i})^{-1} \circ \mathcal{A}_{y_i} - \text{Id}$. Since \mathcal{A}_x is α -Hölder (1.1) and $\|(\mathcal{A}_x)^{-1}\|$ is uniformly bounded we obtain

$$(4.6) \quad \|\Delta_i\| \leq \|(\mathcal{A}_{x_i})^{-1}\| \cdot \|\mathcal{A}_{y_i} - \mathcal{A}_{x_i}\| \leq M' \text{dist}(x_i, y_i)^\alpha \leq M' (\delta e^{-\gamma \min\{i, m-i\}})^\alpha,$$

where the constant M' depends only on the cocycle \mathcal{A} .

Since both x and $f^m x$ are in \mathcal{R}_ℓ we have $K(x_i) \leq \ell e^{\epsilon \min\{i, m-i\}}$ by (3.4) and (4.1). Also, for any points $x, y \in \mathcal{R}$ the inequality (3.3) yields

$$(4.7) \quad \|A\|_{y \leftarrow x} \leq K(y) \|A\| \quad \text{and} \quad \|A\| \leq K(x) \|A\|_{y \leftarrow x}.$$

Using the first inequality we conclude that

$$\begin{aligned} \|\Delta_i\|_{x_i \leftarrow x_i} &\leq K(x_i) \|\Delta_i\| \leq \ell e^{\epsilon \min\{i, m-i\}} \|\Delta_i\| \leq \ell e^{\epsilon \min\{i, m-i\}} M' \delta^\alpha e^{-\gamma \alpha \min\{i, m-i\}} \\ \text{and} \quad \|(\mathcal{A}_{x_i})^{-1} \circ \mathcal{A}_{y_i}\|_{x_i \leftarrow x_i} &\leq 1 + \|\Delta_i\|_{x_i \leftarrow x_i} \leq 1 + M' \ell \delta^\alpha e^{(\epsilon - \alpha\gamma) \min\{i, m-i\}}. \end{aligned}$$

Combining this with (4.5) we obtain

$$\begin{aligned} \log(\|\mathcal{A}_y^n\|_{x_n \leftarrow x_0}) - n(\lambda + \epsilon) &\leq \sum_{i=0}^{n-1} \log(\|(\mathcal{A}_{x_i})^{-1} \circ \mathcal{A}_{y_i}\|_{x_i \leftarrow x_i}) \\ &\leq M' \ell \delta^\alpha \sum_{i=0}^{n-1} e^{(\epsilon - \alpha\gamma) \min\{i, m-i\}} \leq M' \ell \delta^\alpha \cdot 2 \sum_{i=0}^{\infty} e^{(\epsilon - \alpha\gamma)i} = c \ell \delta^\alpha \end{aligned}$$

since $\epsilon < \alpha\gamma$. The constant c depends only on the cocycle \mathcal{A} and on $(\alpha\gamma - \epsilon)$. We conclude using (3.2) that

$$(4.8) \quad \|\mathcal{A}_y^n\|_{x_n \leftarrow x_0} \leq e^{c\ell\delta^\alpha} e^{n(\lambda+\epsilon)}.$$

Since $K(x_0) \leq \ell$ we can also estimate the standard norm using the second inequality in (4.7)

$$\|\mathcal{A}_y^n\| \leq K(x_0) \|\mathcal{A}_y^n\|_{x_n \leftarrow x_0} \leq \ell e^{c\ell\delta^\alpha} e^{n(\lambda+\epsilon)}.$$

The proof of (4.4) is similar. The previous argument with \mathcal{A}_x replaced by $(\mathcal{A}_x)^{-1}$ yields

$$\|(\mathcal{A}_y^n)^{-1}\|_{x_0 \leftarrow x_n} \leq e^{c\ell\delta^\alpha} e^{n(-\chi+\epsilon)}.$$

Then the standard norm can be estimated as $\|(\mathcal{A}_y^n)^{-1}\| \leq K(x_n)\|(\mathcal{A}_y^n)^{-1}\|_{x_0 \leftarrow x_n}$ and (4.4) follows since $K(x_n) \leq \ell e^{\epsilon \min\{n, m-n\}}$. \square

In the proof of the theorem we will also use the following results by A. Karlsson and G. Margulis and by M. Guysinsky.

Proposition 4.2. [KaM99, Proposition 4.2] *Let $a_n(x)$ be an integrable subadditive cocycle with exponent $\lambda > -\infty$ over an ergodic measure-preserving system (X, f, μ) . Then there exists a set $E \subset X$ with $\mu(E) = 1$ such that for each $x \in E$ and each $\epsilon > 0$ there exists an integer $L = L(x, \epsilon)$ and infinitely many n such that*

$$(4.9) \quad a_n(x) - a_{n-i}(f^i x) \geq (\lambda - \epsilon)i \quad \text{for all } i \text{ with } L \leq i \leq n.$$

Lemma 4.3. [G, Lemma 8] *Let $f : X \rightarrow X$ be a homeomorphism preserving an ergodic Borel probability measure μ . Then there exists a set P with $\mu(P) = 1$ such that for each $x \in P$ and $\epsilon, \delta > 0$ there exists an integer $N = N(x, \epsilon, \delta)$ such that if $n > N$ then there is an integer k with*

$$n(1 + \epsilon) < k < n(1 + 2\epsilon) \quad \text{and} \quad \text{dist}(x, f^k x) < \delta.$$

4.2. Finding p such that $|\lambda(\mathcal{A}, \mu) - \frac{1}{k} \log \|\mathcal{A}_p^k\|| < \epsilon$.

We fix $0 < \epsilon < \min\{1, \alpha\gamma/3\}$, the Lyapunov norm $\|\cdot\|_{x, \epsilon}$, the function $K = K_\epsilon$, and the sets \mathcal{R} and \mathcal{R}_ℓ as before. Without loss of generality we can assume that $\mathcal{R} \subset (E \cap P)$, where the set E is given by Proposition 4.2 with $a_n(x) = \log \|\mathcal{A}_x^n\|$ and P is given by Lemma 4.3. We take $\epsilon' = 3\epsilon/(\alpha\gamma)$ and choose ℓ so that $\mu(\mathcal{R}_\ell) > 1 - \epsilon'/2$.

First we describe the choice of the periodic point $p = f^k p$. We fix a point $x \in \mathcal{R}_\ell$ for which the Birkhoff Ergodic Theorem holds for the indicator function of \mathcal{R}_ℓ :

$$(4.10) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \#\{i : 1 \leq i \leq n \text{ and } f^i x \in \mathcal{R}_\ell\} = \mu(\mathcal{R}_\ell) > 1 - \epsilon'/2.$$

We fix $L = L(x, \epsilon)$ given by Proposition 4.2. We then take $\delta > 0$ sufficiently small so that (4.17) is satisfied. We fix $N = N(x, \epsilon', \delta/D)$ given by Lemma 4.3, where D is as in the closing property, Definition 1.3. By Proposition 4.2 there are arbitrarily large n satisfying (4.9). We consider such an n greater than N and L . Then by Lemma 4.3 there exists k such that $n(1 + \epsilon') < k < n(1 + 2\epsilon')$ and $\text{dist}(x, f^k x) < \delta/D$. Then by the closing property there exists a periodic point $p = f^k p$ such that

$$(4.11) \quad \text{dist}(f^i x, f^i p) \leq \delta e^{-\gamma \min\{i, k-i\}} \quad \text{for every } i = 0, \dots, k.$$

By (4.10), if n is sufficiently large then there exists m such that $f^m x \in \mathcal{R}_\ell$ and $n \leq m \leq n(1 + \epsilon') < k$. We summarize our choices:

$$\begin{aligned} \max\{L, N\} < n \leq m \leq n(1 + \epsilon') < k < n(1 + 2\epsilon'), \\ n \text{ satisfies (4.9), } \quad x, f^m x \in \mathcal{R}_\ell, \quad p = f^k p \text{ satisfies (4.11).} \end{aligned}$$

First we obtain an upper estimate for $\|\mathcal{A}_p^k\|$. Since (4.11) also holds with m in place of k , we can apply Lemma 4.1 with $y = p$ to get

$$\|\mathcal{A}_p^m\| \leq \ell e^{c\ell\delta^\alpha} e^{m(\lambda+\epsilon)}.$$

As $k - m < 2\epsilon'n < 2\epsilon'k$, we have

$$\begin{aligned} \|\mathcal{A}_p^k\| &\leq \|\mathcal{A}_p^m\| \cdot \|\mathcal{A}_{f^m p}^{k-m}\| \leq \ell e^{c\ell\delta^\alpha} e^{m(\lambda+\epsilon)} \cdot e^{\lambda'(k-m)} = \ell e^{c\ell\delta^\alpha} e^{k(\lambda+\epsilon)+(\lambda'-\lambda-\epsilon)(k-m)} \\ &\leq \ell e^{c\ell\delta^\alpha} e^{k(\lambda+\epsilon)+(\lambda'-\lambda)2\epsilon'k}. \end{aligned}$$

Taking logarithm we obtain

$$\frac{1}{k} \log \|\mathcal{A}_p^k\| \leq \lambda + \epsilon + (\lambda' - \lambda)2\epsilon' + \frac{1}{k}(\log \ell + c\ell\delta^\alpha).$$

Taking n , and hence k , sufficiently large we conclude that

$$(4.12) \quad \frac{1}{k} \log \|\mathcal{A}_p^k\| \leq \lambda + \epsilon + (\lambda' - \lambda)2\epsilon' + \epsilon = \lambda + 2\epsilon + 6\epsilon(\lambda' - \lambda)/(\alpha\gamma).$$

Now we obtain the lower estimate for $\|\mathcal{A}_p^k\|$. First we bound $\|\mathcal{A}_x^n - \mathcal{A}_p^n\|$.

$$\begin{aligned} \mathcal{A}_x^n - \mathcal{A}_p^n &= \mathcal{A}_{x_1}^{n-1} \circ (\mathcal{A}_x - \mathcal{A}_p) + (\mathcal{A}_{x_1}^{n-1} - \mathcal{A}_{p_1}^{n-1}) \circ \mathcal{A}_p \\ &= \mathcal{A}_{x_1}^{n-1} \circ (\mathcal{A}_x - \mathcal{A}_p) + ((\mathcal{A}_{x_2}^{n-2} \circ (\mathcal{A}_{x_1} - \mathcal{A}_{p_1}) + (\mathcal{A}_{x_2}^{n-2} - \mathcal{A}_{p_2}^{n-2}) \circ \mathcal{A}_{p_1}) \circ \mathcal{A}_p) \\ &= \mathcal{A}_{x_1}^{n-1} \circ (\mathcal{A}_x - \mathcal{A}_p) + \mathcal{A}_{x_2}^{n-2} \circ (\mathcal{A}_{x_1} - \mathcal{A}_{p_1}) \circ \mathcal{A}_p + (\mathcal{A}_{x_2}^{n-2} - \mathcal{A}_{p_2}^{n-2}) \circ \mathcal{A}_p^2 \\ &= \dots = \sum_{i=0}^{n-1} \mathcal{A}_{x_{i+1}}^{n-(i+1)} \circ (\mathcal{A}_{x_i} - \mathcal{A}_{p_i}) \circ \mathcal{A}_p^i. \end{aligned}$$

Hence we can estimate the norm as follows

$$(4.13) \quad \|\mathcal{A}_x^n - \mathcal{A}_p^n\| \leq \sum_{i=0}^{n-1} \|\mathcal{A}_{x_{i+1}}^{n-(i+1)}\| \cdot \|\mathcal{A}_{x_i} - \mathcal{A}_{p_i}\| \cdot \|\mathcal{A}_p^i\|.$$

Since n satisfies (4.9) of Proposition 4.2 with $a_n(x) = \log \|\mathcal{A}_x^n\|$,

$$a_{n-i}(x_i) \leq a_n(x) - (\lambda - \epsilon)i \quad \text{for all } i \text{ with } L \leq i \leq n,$$

and thus for all such i

$$\|\mathcal{A}_{x_{i+1}}^{n-(i+1)}\| \leq \|\mathcal{A}_x^n\| e^{-(i+1)(\lambda-\epsilon)}.$$

Using (4.11) and Hölder continuity of \mathcal{A} we obtain

$$\|\mathcal{A}_{x_i} - \mathcal{A}_{p_i}\| \leq M \text{dist}(x_i, p_i)^\alpha \leq M(\delta e^{-\gamma \min\{i, k-i\}})^\alpha \leq M\delta^\alpha e^{-\alpha\gamma \min\{i, k-i\}}.$$

To estimate the exponent we claim that

$$\alpha\gamma \min\{i, k-i\} \geq 3\epsilon i \quad \text{for } i = 1, \dots, n.$$

Indeed, this holds if $i = \min\{i, k - i\}$ as $\epsilon < \alpha\gamma/3$. If $k - i = \min\{i, k - i\}$ then

$$\alpha\gamma(k - i) \geq 3\epsilon i \iff (3\epsilon + \alpha\gamma)i \leq \alpha\gamma k \iff i \leq \frac{k}{1 + 3\epsilon/(\alpha\gamma)},$$

which holds for $i \leq n$ since $n < k/(1 + \epsilon')$ and $\epsilon' = 3\epsilon/(\alpha\gamma)$. Thus we conclude that

$$(4.14) \quad \|\mathcal{A}_{x_i} - \mathcal{A}_{p_i}\| \leq M\delta^\alpha e^{-3\epsilon i} \quad \text{for all } i = 1, \dots, n.$$

Applying Lemma 4.1 as before, we get $\|\mathcal{A}_y^i\| \leq \ell e^{c\ell\delta^\alpha} e^{i(\lambda+\epsilon)}$, for all $i = 1, \dots, m$. Combining these estimates we obtain that for $L \leq i \leq n$

$$\begin{aligned} & \|\mathcal{A}_{x_{i+1}}^{n-(i+1)}\| \cdot \|\mathcal{A}_{x_i} - \mathcal{A}_{p_i}\| \cdot \|\mathcal{A}_p^i\| \leq \\ & \leq \|\mathcal{A}_x^n\| e^{-(i+1)(\lambda-\epsilon)} \cdot M\delta^\alpha e^{-3\epsilon i} \cdot \ell e^{c\ell\delta^\alpha} e^{i(\lambda+\epsilon)} = C_1(\delta) \|\mathcal{A}_x^n\| e^{-\epsilon i}. \end{aligned}$$

where $C_1(\delta) = \delta^\alpha M \ell e^{c\ell\delta^\alpha - \lambda + \epsilon}$. We conclude that

$$(4.15) \quad \begin{aligned} & \sum_{i=L}^{n-1} \|\mathcal{A}_{x_{i+1}}^{n-(i+1)}\| \cdot \|\mathcal{A}_{x_i} - \mathcal{A}_{p_i}\| \cdot \|\mathcal{A}_p^i\| \\ & \leq C_1(\delta) \|\mathcal{A}_x^n\| \sum_{i=L}^{n-1} e^{-\epsilon i} \leq C_1(\delta) \|\mathcal{A}_x^n\| \frac{1}{1 - e^{-\epsilon}} = C_2(\delta) \|\mathcal{A}_x^n\|. \end{aligned}$$

where $C_2(\delta) = C_1(\delta)(1 - e^{-\epsilon})^{-1}$.

For $i < L$ we estimate $\|\mathcal{A}_{x_i}^{n-i}\| \leq \|\mathcal{A}_x^n\| \cdot \|(\mathcal{A}_x^i)^{-1}\| \leq \|\mathcal{A}_x^n\| e^{-\chi' i}$, where $\chi' < \chi$ is such that $\|(\mathcal{A}_x)^{-1}\| \leq e^{-\chi'}$ for all $x \in X$. Hence

$$(4.16) \quad \begin{aligned} & \sum_{i=0}^{L-1} \|\mathcal{A}_{x_{i+1}}^{n-(i+1)}\| \cdot \|\mathcal{A}_{x_i} - \mathcal{A}_{p_i}\| \cdot \|\mathcal{A}_p^i\| \\ & \leq \sum_{i=0}^{L-1} \|\mathcal{A}_x^n\| e^{-(i+1)\chi'} \cdot M\delta^\alpha e^{-3\epsilon i} \cdot \ell e^{c\ell\delta^\alpha} e^{i(\lambda+\epsilon)} \leq C_3(\delta) \|\mathcal{A}_x^n\|, \end{aligned}$$

where $C_3(\delta) = L e^{-\chi'+(\lambda-\chi')L} \delta^\alpha M \ell e^{c\ell\delta^\alpha}$, as $\lambda - \chi' > 0$.

Combining estimates (4.13), (4.15) and (4.16) we obtain

$$\|\mathcal{A}_x^n - \mathcal{A}_p^n\| \leq \|\mathcal{A}_x^n\| (C_2(\delta) + C_3(\delta)) \leq \|\mathcal{A}_x^n\|/2$$

since by the choice of $\delta > 0$ we have

$$(4.17) \quad C_2(\delta) + C_3(\delta) = \delta^\alpha M \ell e^{c\ell\delta^\alpha} \left((1 - e^{-\epsilon})^{-1} e^{-\lambda+\epsilon} + L e^{-\chi'+(\lambda-\chi')L} \right) < 1/2.$$

Hence

$$\|\mathcal{A}_p^n\| \geq \|\mathcal{A}_x^n\| - \|\mathcal{A}_x^n - \mathcal{A}_p^n\| \geq \|\mathcal{A}_x^n\|/2 > e^{(\lambda-\epsilon)n}/2,$$

provided that n is sufficiently large for the limit in (2.3). Since $\mathcal{A}_p^n = (\mathcal{A}_{f_p^{n-p}}^{k-n})^{-1} \circ \mathcal{A}_p^k$,

$$\|\mathcal{A}_p^n\| \leq \|(\mathcal{A}_{f_p^{n-p}}^{k-n})^{-1}\| \cdot \|\mathcal{A}_p^k\| \leq \|\mathcal{A}_p^k\| e^{-\chi'(k-n)}.$$

Hence

$$\|\mathcal{A}_p^k\| \geq e^{\chi'(k-n)} \|\mathcal{A}_p^n\| > e^{(\lambda-\epsilon)n + \chi'(k-n)}/2 > e^{(\lambda-\epsilon)k - (\lambda-\chi')(k-n)}/2,$$

$$\text{and so } \frac{1}{k} \log \|\mathcal{A}_p^k\| > \frac{1}{k} [(\lambda - \epsilon)k - (\lambda - \chi')(k - n) - \log 2].$$

Since $k - n < 2\epsilon'n < 2\epsilon'k$ we obtain

$$\frac{1}{k} \log \|\mathcal{A}_p^k\| > (\lambda - \epsilon) - (\lambda - \chi')2\epsilon' - \log 2/k > \lambda - 2\epsilon - 6\epsilon(\lambda - \chi')/(\alpha\gamma)$$

if n and hence k are sufficiently large.

We conclude that for each $\epsilon > 0$ there exists $p = f^k p$ satisfying this equation as well as (4.12), and the approximation of $\lambda(\mathcal{A}, \mu)$ by $\frac{1}{k} \log \|\mathcal{A}_p^k\|$ follows.

4.3. Finding p approximating both $\lambda(\mathcal{A}, \mu)$ and $\chi(\mathcal{A}, \mu)$.

We describe how to modify the previous argument to obtain p approximating both the upper and the lower exponents. The construction of p and the calculations are similar. The main difference is that for the point x we need to find arbitrarily large n satisfying both (4.9) and

$$(4.18) \quad \tilde{a}_n(x) - \tilde{a}_{n-i}(f^i x) \geq (-\chi - \epsilon)i \quad \text{for all } i \text{ with } L \leq i \leq n,$$

where $\tilde{a}_n(x)$ is from (2.2). For this we use an advanced version of Proposition 4.2 due to S. Gouëzel and A. Karlsson.

Proposition 4.4. [GoKa, Theorem 1.1 and Remark 1.2] *Let $a_n(x)$ be an integrable subadditive cocycle with exponent $\lambda > -\infty$ over an ergodic measure-preserving system (X, f, μ) . Then for each $\rho > 0$ there exists a sequence $\epsilon_i \rightarrow 0$ and a set $E_\rho \subset X$ with $\mu(E_\rho) > 1 - \rho$ such that for each $x \in E_\rho$ the set S of integers n satisfying*

$$(4.19) \quad a_n(x) - a_{n-i}(f^i x) \geq (\lambda - \epsilon_i)i \quad \text{for all } i \text{ with } 1 \leq i \leq n$$

has asymptotic upper density is greater than $1 - \rho$, that is

$$\overline{\text{Dens}}(S) \stackrel{\text{def}}{=} \limsup \frac{1}{N} |S \cap [0, N - 1]| > 1 - \rho.$$

We take $\rho < 1/2$ and use Proposition 4.4 to obtain sets E_ρ and \tilde{E}_ρ of measure greater than $1 - \rho$ for $a_n(x)$ and $\tilde{a}_n(x)$ respectively. As in the previous argument, we take $x \in \mathcal{R}_\ell$ for which Birkhoff Ergodic Theorem holds for the indicator function of \mathcal{R}_ℓ and, in addition, require that $x \in E_\rho \cap \tilde{E}_\rho$. Then Proposition 4.4 ensures that there exist infinitely many n satisfying (4.19) simultaneously for $a_n(x)$ and for $\tilde{a}_n(x)$ with exponent χ . Moreover, there exists $L = L(\epsilon)$ so that $\tilde{\epsilon}_i, \epsilon_i < \epsilon$ for all $i \geq L$ and thus we obtain that for such n we have both (4.9) and (4.18).

All other choices remain the same except we take $\epsilon' = 4\epsilon/(\alpha\gamma)$. The argument for λ is unchanged. The upper estimate for $-\chi$ is the same direct application of (4.4) in

Lemma 4.1. To obtain the lower estimate of $-\chi$ we use the equation

$$(\mathcal{A}_x^n)^{-1} - (\mathcal{A}_p^n)^{-1} = \sum_{i=0}^{n-1} (\mathcal{A}_p^i)^{-1} \circ ((\mathcal{A}_{x_i})^{-1} - (\mathcal{A}_{p_i})^{-1}) \circ (\mathcal{A}_{x_{i+1}}^{n-(i+1)})^{-1},$$

which yields an inequality for the norm similar to (4.13). The first and third terms are then estimated using (4.4) and (4.18). We use the new choice of ϵ' to get $e^{-4\epsilon i}$ decay in (4.14) to compensate for the extra term in (4.4) compared to (4.3). \square

5. PROOFS OF PROPOSITION 1.5 AND COROLLARY 1.6

Let μ_p be the uniform measure on the orbit of a periodic point $p = f^k p$. Then

$$\begin{aligned} \lambda(\mathcal{A}, \mu_p) &= \lim_{N \rightarrow \infty} \frac{1}{N} \log \|\mathcal{A}_p^N\| = \lim_{n \rightarrow \infty} \frac{1}{nk} \log \|(\mathcal{A}_p^k)^n\| = \\ &= \frac{1}{k} \log \left(\lim_{n \rightarrow \infty} \|(\mathcal{A}_p^k)^n\|^{1/n} \right) = \frac{1}{k} \log (r(\mathcal{A}_p^k)) \leq \frac{1}{k} \log \|\mathcal{A}_p^k\|, \end{aligned}$$

where $r(A) = \lim_{n \rightarrow \infty} \|A^n\|^{1/n}$ is the spectral radius of a linear operator A . Similarly,

$$-\chi(\mathcal{A}, \mu_p) = \frac{1}{k} \log (r((\mathcal{A}_p^k)^{-1})), \quad \text{so} \quad \chi(\mathcal{A}, \mu_p) \geq \frac{1}{k} \log \|(\mathcal{A}_p^k)^{-1}\|^{-1}.$$

It follows that for p as in Theorem 1.4 we have the one-sided estimates (1.6).

5.1. Proof of Proposition 1.5. Suppose that f is a homeomorphisms of a compact metric space X and $a_n(x)$ is a continuous subadditive cocycle. Then it is easy to see that $a_n = \sup_{x \in X} a_n(x)$ is a subadditive sequence and so there is limit

$$\hat{\nu}(a) := \lim_{n \rightarrow \infty} a_n/n, \quad \text{where} \quad a_n = \sup_{x \in X} a_n(x).$$

By Theorem 1 in [Sch98], $\hat{\nu}(a) = \sup_{\mu} \nu(a, \mu)$ where supremum is taken over all f -invariant ergodic probability measures on X .

Let $a_n(x) = \log \|\mathcal{A}_x^n\|$, where \mathcal{A} is a Hölder continuous Banach cocycle over f , and let f be a homeomorphism of a compact metric space X satisfying closing property. We denote $\hat{\lambda}(\mathcal{A}) = \sup_{\mu} \lambda(\mathcal{A}, \mu)$. Then we have

$$\hat{\lambda}(\mathcal{A}) := \sup_{\mu} \lambda(\mathcal{A}, \mu) = \sup_{\mu} \nu(a, \mu) = \hat{\nu}(a).$$

We also consider the supremum over all periodic measures μ_p and denote

$$\hat{\lambda}_p(\mathcal{A}) := \sup_{\mu_p} \lambda(\mathcal{A}, \mu_p) = \sup \left\{ \frac{1}{k} \log (r(\mathcal{A}_p^k)) : p = f^k p \right\}.$$

If V is finite dimensional [K11, Theorem 1.4] yields that $\hat{\lambda}(\mathcal{A}) = \hat{\lambda}_p(\mathcal{A})$. However, for infinite dimensional space $\hat{\lambda}_p(\mathcal{A})$ can be smaller than $\hat{\lambda}(\mathcal{A})$.

Indeed, let $f : X \rightarrow X$ be the full shift on two symbols, i.e.

$$X = \{ \bar{x} = (x_n)_{n \in \mathbb{Z}} : x_n \in \{0, 1\} \} \quad \text{and} \quad f(\bar{x}) = (x_{n+1})_{n \in \mathbb{Z}}$$

and let \mathcal{A} be a cocycle with values in bounded operators on Hilbert space ℓ_2 given by

$$\mathcal{A}_{\bar{x}} = B_0 \text{ if } x_0 = 0 \quad \text{and} \quad \mathcal{A}_{\bar{x}} = B_1 \text{ if } x_0 = 1.$$

Since any \mathcal{A}_x^n is a product of B_0 and B_1 of length n , and any such product is \mathcal{A}_x^n for some $x \in X$, we have

$$\begin{aligned} e^{\hat{\lambda}(\mathcal{A})} &= \lim_{n \rightarrow \infty} \sup \{ e^{a_n(x)/n} : x \in X \} = \lim_{n \rightarrow \infty} \sup \{ \|\mathcal{A}_x^n\|^{1/n} : x \in X \} \\ &= \lim_{n \rightarrow \infty} \sup \{ \|A_n \cdots A_1\|^{1/n} : A_i \in \{B_0, B_1\} \text{ for } i = 1, \dots, n \} =: \hat{\rho}(B_0, B_1). \end{aligned}$$

Also, as any product of B_0 and B_1 of length n can be realized as \mathcal{A}_p^n for some $p = f^n p$

$$\begin{aligned} e^{\hat{\lambda}_p(\mathcal{A})} &= \sup \left\{ (r(\mathcal{A}_p^k))^{1/k} : p = f^k p \right\} = \\ &= \sup \left\{ (r(A_k \cdots A_1))^{1/k} : k \in \mathbb{N}, A_i \in \{B_0, B_1\} \right\} = \\ &\stackrel{(*)}{=} \limsup_{k \rightarrow \infty} \sup \left\{ (r(A_k \cdots A_1))^{1/k} : A_i \in \{B_0, B_1\} \text{ for } i = 1, \dots, k \right\} =: \bar{\rho}(B_0, B_1). \end{aligned}$$

Equality $(*)$ holds since $r(A) = r(A^m)^{1/m}$ and so taking $A_k \cdots A_1$ with $r(A_k \cdots A_1)^{1/k}$ close to the supremum and repeating it we obtain an arbitrarily long product with the same value. The number $\hat{\rho}(B_0, B_1)$ is called the *joint spectral radius* of B_0 and B_1 , and $\bar{\rho}(B_0, B_1)$ is called the *generalized spectral radius*.

By Theorem A.1 in [Gu], for any $0 < \alpha < \beta$ there exist two isomorphisms B_0 and B_1 of ℓ_2 such that $\bar{\rho}(B_0, B_1) = \alpha$ and $\hat{\rho}(B_0, B_1) = \beta$. Hence for the corresponding cocycle \mathcal{A} we have $\hat{\lambda}_p(\mathcal{A}) < \hat{\lambda}(\mathcal{A})$. It follows that there is an ergodic measure μ so that $\lambda(\mathcal{A}, \mu) > \sup_{\mu_p} \lambda(\mathcal{A}, \mu_p)$. \square

5.2. Proof of Corollary 1.6. We prove the second part, and the first one is obtained similarly. We denote

$$\begin{aligned} \hat{\sigma}(\mathcal{A}) &= \lim_{n \rightarrow \infty} \sup \{ Q(x, n)^{1/n} : x \in X \}, \\ \hat{\sigma}_p(\mathcal{A}) &= \limsup_{k \rightarrow \infty} \sup \{ Q(p, k)^{1/k} : p = f^k p \}, \\ q_n(x) &= \log Q(x, n) = \log \|\mathcal{A}_x^n\| + \log \|(\mathcal{A}_x^n)^{-1}\|. \end{aligned}$$

Since $a_n(x) = \log \|\mathcal{A}_x^n\|$ and $\tilde{a}_n(x) = \log \|(\mathcal{A}_x^n)^{-1}\|$ are subadditive cocycles over f , so is $q_n(x) = a_n(x) + \tilde{a}_n(x)$. For its exponent $\nu(q, \mu)$ we have

$$\nu(q, \mu) = \nu(a, \mu) + \nu(\tilde{a}, \mu) = \lambda(\mathcal{A}, \mu) - \chi(\mathcal{A}, \mu).$$

As in the proof of Proposition 1.5 we consider

$$\hat{\nu}(q) = \sup_{\mu} \nu(q, \mu) = \lim_{n \rightarrow \infty} q_n/n, \quad \text{where } q_n = \sup_{x \in X} q_n(x) = \log \sup_{x \in X} Q(x, n).$$

It follows from Theorem 1.4 that for any $N \in \mathbb{N}$

$$\begin{aligned} \hat{\nu}(q) &= \sup_{\mu} \nu(q, \mu) = \sup_{\mu} (\lambda(\mathcal{A}, \mu) - \chi(\mathcal{A}, \mu)) \leq \\ &\leq \sup \left\{ \frac{1}{k} \log \|\mathcal{A}_p^k\| + \frac{1}{k} \log \|(\mathcal{A}_p^k)^{-1}\| : p = f^k p, k \geq N \right\} = \\ &= \sup \left\{ \frac{1}{k} \log Q(p, k) : p = f^k p, k \geq N \right\}. \end{aligned}$$

Since $\log \hat{\sigma}(\mathcal{A}) = \hat{\nu}(q)$ it follows that $\hat{\sigma}(\mathcal{A}) \leq \hat{\sigma}_p(\mathcal{A})$. The opposite inequality is clear, and so $\hat{\sigma}(\mathcal{A}) = \hat{\sigma}_p(\mathcal{A})$.

Suppose that $Q(p, k) \leq Ce^{sk}$ whenever $p = f^k p$. Then we have

$$s \geq \log \hat{\sigma}_p(\mathcal{A}) = \log \hat{\sigma}(\mathcal{A}) = \hat{\nu}(q) = \lim_{n \rightarrow \infty} q_n/n.$$

It follows that for each $\epsilon > 0$ there exists $N \in \mathbb{N}$ such that $q_n \leq (s + \epsilon)n$ for all $n > N$ and hence $Q(x, n) \leq e^{(s+\epsilon)n}$ for all $x \in X$ and $n > N$. Taking

$$C_\epsilon = \max \{ Q(x, n) : x \in X \text{ and } 1 \leq n \leq N \},$$

we obtain $Q(x, n) \leq C_\epsilon e^{(s+\epsilon)n}$ for all $x \in X$ and $n \in \mathbb{N}$.

The statements for negative n follow as $Q(x, -n) = Q(f^{-n}x, n)$. Hence

$$\sup \{ Q(x, -n)^{1/|n|} : x \in X \} = \sup \{ Q(x, n)^{1/|n|} : x \in X \}$$

and so the limit as $n \rightarrow -\infty$ equals $\hat{\sigma}(\mathcal{A})$. \square

REFERENCES

- [BP] L. Barreira and Ya. Pesin. *Nonuniformly hyperbolicity: dynamics of systems with nonzero Lyapunov exponents*. Encyclopedia of Mathematics and Its Applications **115**, Cambridge University Press, 2007.
- [Gu] L. Gurvits. *Stability of discrete linear inclusion*. Linear Algebra and its Applications, vol. 231 (1995), 47-85.
- [G] M. Guysinsky. *Livšic theorem for Banach rings*. Preprint.
- [GoKa] S. Gouëzel and A. Karlsson. *Subadditive and multiplicative ergodic theorems*. Preprint.
- [K11] B. Kalinin. *Livsic Theorem for matrix cocycles*. Annals of Math. **173** (2011), no. 2, 1025-1042.
- [KS13] B. Kalinin and V. Sadovskaya. *Cocycles with one exponent over partially hyperbolic systems*. Geometriae Dedicata, Vol. 167, Issue 1 (2013), 167-188.
- [KaM99] A. Karlsson and G. Margulis. *A multiplicative ergodic theorem and nonpositively curved spaces*. Communications in Mathematical Physics **208** (1999) 107-123.
- [KtH] A. Katok and B. Hasselblatt. *Introduction to the modern theory of dynamical systems*. Encyclopedia of Mathematics and its Applications **54**, Cambridge University Press, 1995.
- [LL] Z. Lian and K. Lu. *Lyapunov exponents and invariant manifolds for random dynamical systems in a Banach space*. Mem. Amer. Math. Soc. **206** (2010), no. 967.
- [LY12] Z. Lian and L. S. Young. *Lyapunov exponents, periodic orbits, and horseshoes for semiflows on Hilbert spaces*. J. Amer. Math. Soc. **25** (2012), no. 3, 637-665.

- [M12] I. Morris. *The generalised Berger-Wang formula and the spectral radius of linear cocycles*. Journal of Functional Analysis **262** (2012) 811-824.
- [Sch98] S. J. Schreiber. *On growth rates of subadditive functions for semi-flows*. J. Differential Equations, 148, 334350, 1998.

DEPARTMENT OF MATHEMATICS, THE PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK,
PA 16802, USA.

E-mail address: kalinin@psu.edu, sadovskaya@psu.edu